

On Pointed Hopf Algebras and Kaplansky's 10th Conjecture

Shlomo Gelaki

Department of Mathematics

Harvard University

Cambridge, MA 02138

March 26, 1998

Radford constructed two families of finite dimensional pointed Hopf algebras over a field k [R1, 5.1 and 5.2]. The first one, denoted by $H_{n,q,N,\nu}$, is a family of pointed Hopf algebras which contains a sub-family of self dual pointed Hopf algebras, denoted by $H_{(N,\nu,\omega)}$. This sub-family generalizes Taft's well known Hopf algebras [G1,T]. The second one, denoted by $U_{(N,\nu,\omega)}$, is a family of pointed, unimodular and ribbon Hopf algebras constructed for the purpose of computing invariants of knots, links and 3-manifolds. This family generalizes the well known quantum group $U_q(sl_2)'$ when q is a root of unity. The author constructed a new family of finite dimensional Hopf algebras, denoted by $\mathcal{H}_{n,q,N,\nu,\alpha}$, which generalizes Radford's $H_{n,q,N,\nu}$, and furthermore proved that this new family captures all Hopf algebras generated as algebras by one grouplike element of order N , and one non-trivial skew primitive element [G3, Theorem 1.1.1]. He also proved that $U_q(sl_2)'$ and the 8-dimensional Hopf algebra $U_{(2,1,-1)}$ capture all minimal quasitriangular Hopf algebras U , generated by one grouplike element of *prime* order N , and two non-trivial $kG(U)$ independent skew primitive elements [G3, Theorem 1.2.2]. The purpose of this paper is three fold. First, to study the family $\mathcal{H}_{n,q,N,\nu,\alpha}$, and then to construct and study a new family of finite dimensional pointed and unimodular Hopf algebras which generalizes Radford's $U_{(N,\nu,\omega)}$. Second, **to answer in the negative Kaplansky's 10th conjecture from 1975** on the finite number of Hopf algebras of a given dimension [K]. Third, to generalize [G3, Theorem 1.2.2] and characterize the sub-family $U_{(N,\nu,\omega)}$ via minimal quasitriangularity.

The paper is organized as follows. In Section 1 we discuss some aspects of the theory of finite dimensional Hopf algebras, and introduce all the Hopf algebras which are relevant for this paper.

In Section 2 we study the family of Hopf algebras $\mathcal{H}_{n,q,N,\nu,\alpha}$ and its family of dual Hopf algebras $\mathcal{H}_{n,q,N,\nu,\alpha}^*$. First, in **Proposition 2.1** we show that $\mathcal{H}_{n,q,N,\nu,\alpha}$ is pointed, find a linear basis for it, find the first term of its coradical filtration, find its left and right integrals and show that it is an extension of $k\mathbb{Z}_{N/n}$ by $H_{n,q,n,\nu}$. Second, in **Lemma 2.2** we describe

$G(\mathcal{H}_{n,q,N,\nu,\alpha}^*)$ (where $G(H)$ denotes the group of grouplike elements of a Hopf algebra H), and then use it in **Theorem 2.4** to show that Radford's family $H_{(N,\nu,\omega)}$ is characterized via self duality. Third, in **Theorem 2.5** we find a necessary and sufficient condition for the dual Hopf algebra $\mathcal{H}_{n,q,N,\nu,\alpha}^*$ to be pointed by constructing irreducible representations of $\mathcal{H}_{n,q,N,\nu,\alpha}$. Forth, in **Proposition 2.6** we prove that Radford's family $H_{n,q,N,\nu}$ is closed under duality, and then use it in **Corollary 2.7** to show that $\mathcal{H}_{n,q,N,\nu,\alpha}^*$ is an extension of $H_{n,q,n,\nu}^* \cong H_{\frac{n}{(n,\nu)},\omega^\nu,N,\mu}$ (hence of $H_{n,q,n,\nu}$ if $\alpha \neq 0$) by $k\mathbb{Z}_{N/n}$. Finally, in **Lemma 2.8** we find all the Hopf subalgebras of $H_{n,q,N,\nu}$, and then use it and Theorem 2.5 to determine in **Theorem 2.9** when $\mathcal{H}_{n,q,N,\nu,\alpha}$ is quasitriangular.

In Section 3 we first construct and study a new family of finite dimensional pointed Hopf algebras, denoted by $\mathcal{U}_{(n,N,\nu,q,\alpha,\beta,\gamma)}$. In particular in **Proposition 3.2** we show that it is pointed, find a linear basis for it, find the first term of its coradical filtration, show that it is an extension of $k\mathbb{Z}_{N/n}$ by $U_{(n,\nu,\omega)}$, for some primitive n th root of unity ω , and prove that it is always unimodular by computing a non-zero 2-sided integral for it. We also show that over any infinite field which contains a primitive n th root of unity, our new family contains infinitely many non-isomorphic Hopf algebras of any dimension of the form Nn^2 , where $2 < n < N$ are integers so that n divides N . Thus, in **Corollary 3.3** we prove that Kaplansky's 10th conjecture from 1975 [K] is false. Second, in **Theorem 3.6** we use Theorem 2.5 to generalize [G3, Theorem 1.2.2], and show that over an algebraically closed field of characteristic zero, Radford's family $U_{(N,\nu,\omega)}$, which we identify as the subfamily $\mathcal{U}_{(\frac{N}{(N,\nu)},N,\nu,\omega^\nu,0,0,1)}$, captures all minimal quasitriangular Hopf algebras U , generated as algebras by one grouplike element of an arbitrary order N , and two non-trivial $kG(U)$ independent skew primitive elements.

I wish to thank the referee for his/her very useful comments.

1 Preliminaries

Throughout this paper, unless otherwise explicitly stated, k denotes a field. Also, we shall denote the greatest common divisor of two integers a and b by (a, b) .

The Hopf algebras which are studied in this paper are pointed. Recall that a Hopf algebra H over the field k is pointed if its simple subcoalgebras are 1-dimensional; that is, they are generated by elements of $G(H)$. For any $g, h \in G(H)$ we denote the vector space of $g : h$ skew primitives of H by $P_{g,h}(H) = \{x \in H \mid \Delta(x) = x \otimes g + h \otimes x\}$. Thus the classical primitive elements are $P_{1,1}(H)$. The element $g - h$ is always $g : h$ skew primitive. If $P_{g,h}(H)$ consists only of the linear span of $g - h$ we say it is trivial. Otherwise, let $P_{g,h}(H)'$ denote a complement of $sp_k\{g - h\}$ in $P_{g,h}(H)$. Taft-Wilson theorem [TW] states that the first term H_1 of the coradical filtration of H is given by:

$$H_1 = kG(H) \bigoplus \left(\bigoplus_{g,h \in G(H)} P_{g,h}(H)' \right). \quad (1)$$

The following lemma is very useful in determining whether certain Hopf algebras are pointed. Various forms of it appear in the literature; see [R1, Lemma 1] for example.

Lemma 1.1 *Suppose that H is a Hopf algebra over the field k which is generated as an algebra by a subset S of $G(H)$ and by $g : g'$ skew primitives, where g, g' run over S . Then H is pointed and $G(H)$ is generated as a group by S .*

The next proposition is very useful, in particular in the context of quantum groups; see [R1, Proposition 1] for example.

Proposition 1.2 *Suppose that A is an algebra over the field k and $\omega \in k$ is a primitive n th root of unity. Let $a, x \in A$ satisfy $xa = \omega ax$. Then $(a + x)^n = a^n + x^n$.*

Lemma 1.3 [G2, Lemma 0.2] *Let H be a finite dimensional Hopf algebra over a field k . Assume that $(\dim H)1 \neq 0$ and let $x \in P_{a,1}(H)$. Then $xa = ax$ if and only if $x \in \text{sp}_k\{a - 1\}$.*

Next, we give a very brief description of the underlying Hopf algebra structure of $D(H)$, the Drinfel'd double of a finite dimensional Hopf algebra H [D]. As a coalgebra $D(H) = H^{*cop} \otimes H$ (where $H^{*cop} = H^*$ as an algebra, with new comultiplication $\Delta^{cop} = \tau \circ \Delta$, where $\tau : H^* \otimes H^* \rightarrow H^* \otimes H^*$ is the usual twist map). Set $p \bowtie a = p \otimes a$ for $p \in H^*$ and $a \in H$. Multiplication in the double can be described in several ways. The following are very convenient:

$$(p \bowtie a)(q \bowtie b) = \sum p(a_{(1)} \cdot q \cdot s^{-1}(a_{(3)})) \bowtie a_{(2)}b$$

and

$$(p \bowtie a)(q \bowtie b) = \sum pq_{(2)} \bowtie ((s^*)^{-1}(q_{(1)}) \rightharpoonup a \leftharpoonup q_{(3)})b$$

where $\langle a \cdot q, b \rangle = \langle q, ba \rangle$, $\langle q \cdot a, b \rangle = \langle q, ab \rangle$, $p \rightharpoonup a = \sum a_{(1)} \langle p, a_{(2)} \rangle$ and $a \leftharpoonup p = \sum \langle p, a_{(1)} \rangle a_{(2)}$ for all $p, q \in H^*$ and $a, b \in H$. The antipode S of $D(H)$ is given by $S(p \bowtie a) = (\varepsilon \bowtie s(a))((s^*)^{-1}(p) \bowtie 1)$ for $p \in H^*$ and $a \in H$. Observe that $(p \bowtie 1)(q \bowtie b) = pq \bowtie b$ and $(p \bowtie a)(\varepsilon \bowtie b) = p \bowtie ab$. As a consequence, the one-one maps $i_H : H \rightarrow D(H)$ and $i_{H^*} : H^{*cop} \rightarrow D(H)$ defined by $i_H(h) = \varepsilon \bowtie h$ and $i_{H^*}(p) = p \bowtie 1$ respectively for $h \in H$ and $p \in H^*$ are Hopf algebra maps.

We recall now the definition of a quasitriangular Hopf algebra. Let H be a Hopf algebra and $R = \sum R^{(1)} \otimes R^{(2)} \in H \otimes H$. The pair (H, R) is said to be a quasitriangular Hopf algebra if the following axioms hold ($r = R$):

$$(QT.1) \quad \sum \Delta(R^{(1)}) \otimes R^{(2)} = \sum R^{(1)} \otimes r^{(1)} \otimes R^{(2)}r^{(2)},$$

$$(QT.2) \quad \sum \varepsilon(R^{(1)})R^{(2)} = 1,$$

$$(QT.3) \quad \sum R^{(1)} \otimes \Delta^{cop}(R^{(2)}) = \sum R^{(1)}r^{(1)} \otimes R^{(2)} \otimes r^{(2)},$$

$$(QT.4) \quad \sum R^{(1)}\varepsilon(R^{(2)}) = 1 \quad \text{and}$$

$$(QT.5) \quad (\Delta^{cop}(h))R = R(\Delta(h)) \quad \text{for all } h \in H.$$

Drinfel'd has shown that the double of any finite dimensional Hopf algebra is quasitriangular [D]. Suppose that $f : H \rightarrow H'$ is a surjective Hopf algebra map and set $R' = (f \otimes f)(R)$. Then (H', R') is quasitriangular. Let (H, R) be quasitriangular and write $R = \sum R^{(1)} \otimes R^{(2)}$ in the shortest possible way. Set $l(R) = \text{sp}_k \{R^{(1)}\}$ and $r(R) = \text{sp}_k \{R^{(2)}\}$. Let H_R be the Hopf subalgebra of H generated by $l(R)$ and $r(R)$. Then $l(R)$ and $r(R)$ are finite dimensional Hopf subalgebras of H , and there exists an isomorphism of Hopf algebras $f : l(R)^{\text{cop}} \rightarrow r(R)$ and a unique surjection of Hopf algebras $F : D(l(R)) \rightarrow H_R$ which satisfies $F|_{l(R)} = i(l(R))$ and $F|_{l(R)^{\text{cop}}} = f$ [R2]. If $H = H_R$ then (H, R) is called a *minimal* quasitriangular Hopf algebra. We shall also say that H is a (minimal) quasitriangular Hopf algebra if there exists $R \in H \otimes H$ such that (H, R) is a (minimal) quasitriangular Hopf algebra.

We now recall the definition of $H_{n,q,N,\nu}$ and $H_{(N,\nu,\omega)}$ [R1, 5.1]. Let n, ν and N be positive integers such that $n|N$ and $1 \leq \nu < N$. Suppose $q \in k$ is a primitive n th root of unity and let $r = |q^\nu|$. As an algebra $H = H_{n,q,N,\nu}$ is generated by a and x which satisfy the relations

$$a^N = 1, \quad x^r = 0 \quad \text{and} \quad xa = qax.$$

The coalgebra structure of H is determined by

$$\Delta(a) = a \otimes a, \quad \Delta(x) = x \otimes a^\nu + 1 \otimes x, \quad \varepsilon(a) = 1 \quad \text{and} \quad \varepsilon(x) = 0.$$

The antipode of H is determined by

$$s(a) = a^{-1} \quad \text{and} \quad s(x) = -q^{-\nu} a^{-\nu} x.$$

By [R1, Proposition 8 and Theorem 4 a)] and Lemma 1.1, H satisfies the following:

(H.1) As an algebra H is generated by a grouplike element a of order N , and by the space $P_{a^\nu,1}(H)$ of dimension ≤ 2 .

(H.2) H is pointed with $G(H) = (a)$, and if $r > 1$ then it is self dual if and only if there exists a primitive N th root of unity $\omega \in k$ such that $q = \omega^\nu$.

If the field k contains a primitive N th root of unity ω such that $q = \omega^\nu$ and $N \nmid \nu^2$, then $H_{n,q,N,\nu}$ is denoted by $H_{(N,\nu,\omega)}$.

We next recall the definition of the family $\mathcal{H}_{n,q,N,\nu,\alpha}$ [G3, 1.1]. Let n, ν and N be positive integers such that $n|N$ and $1 \leq \nu < N$, and let $\alpha \in k$. Suppose $q \in k$ is a primitive n th root of unity and let $r = |q^\nu|$. For $\alpha = 0$ we define $\mathcal{H}_0 = \mathcal{H}_{n,q,N,\nu,0} = H_{n,q,N,\nu}$. For $0 \neq \alpha \in k$ we further assume that $(n, \nu) = 1$ (hence $r = n$), $n > 1$ and $N \nmid \nu n$. In this case, as an algebra $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$ is generated by a and x which satisfy the relations

$$a^N = 1, \quad x^n = \alpha(a^{\nu n} - 1) \quad \text{and} \quad xa = qax.$$

The coalgebra structure of $\mathcal{H}_\alpha$ is determined by

$$\Delta(a) = a \otimes a, \quad \Delta(x) = x \otimes a^\nu + 1 \otimes x, \quad \varepsilon(a) = 1 \quad \text{and} \quad \varepsilon(x) = 0.$$

The antipode of $\mathcal{H}_\alpha$ is determined by

$$s(a) = a^{-1} \quad \text{and} \quad s(x) = -q^{-\nu} a^{-\nu} x.$$

Note that since $N \nmid \nu n$ it follows that $x^n \neq 0$.

Remark 1.4 Suppose $\alpha \neq 0$. Then to construct $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$ we follow the line of argument and notations used by Radford in order to construct $H_{n,q,N,\nu}$ [R1, 5.1]. We let C be the 4-dimensional coalgebra over k with basis $\{A, B, D, X\}$ whose structure is determined by

$$\Delta(A) = A \otimes A, \quad \Delta(B) = B \otimes B, \quad \Delta(D) = D \otimes D \quad \text{and} \quad \Delta(X) = X \otimes B + D \otimes X, \quad (2)$$

and let I be the ideal of the tensor algebra $T(C)$ generated by

$$A^N - 1, \quad B - A^\nu, \quad D - 1, \quad XA - qAX \quad \text{and} \quad X^n - \alpha(A^{\nu n} - 1).$$

It is straightforward to check that $T(C)$ has a unique bialgebra structure determined by (2), that I is a biideal of $T(C)$ and finally that $\mathcal{H}_\alpha = T(C)/I$, where $a = \overline{A}$ and $x = \overline{X}$, is a Hopf algebra.

In the following we state that over an algebraically closed field of characteristic zero, the family $\mathcal{H}_{n,q,N,\nu,\alpha}$ is unique with respect to property (H.1). We also prove it here for the sake of completeness.

Theorem 1.5 [G3, Theorem 1.1.1] *Let k be an algebraically closed field of characteristic zero. Let $\mathcal{H}$ be a Hopf algebra over k , and suppose there exist integers $1 \leq \nu < N$ so that as an algebra $\mathcal{H}$ is generated by a grouplike element a of order N , and by the space $P_{a^\nu,1}(\mathcal{H})$ of dimension ≤ 2 . Then there exist a positive integer n such that $n|N$, $\alpha \in k$ and a primitive n th root of unity $q \in k$ such that $\mathcal{H} = \mathcal{H}_{n,q,N,\nu,\alpha}$.*

Proof: If $\dim_k P_{a^\nu,1}(\mathcal{H}) = 1$ then $P_{a^\nu,1}(\mathcal{H}) = \text{sp}_k\{a^\nu - 1\}$. Thus, $\mathcal{H} = k(a) = \mathcal{H}_{1,1,N,\nu,0}$, and we are done. Now assume $\dim_k P_{a^\nu,1}(\mathcal{H}) = 2$. First note that the semisimple commutative algebra $k(a)$ acts on $P_{a^\nu,1}(\mathcal{H})$ by conjugation and that $\text{sp}_k\{a^\nu - 1\}$ is a $k(a)$ -submodule of $P_{a^\nu,1}(\mathcal{H})$. Thus there exists a 1-dimensional $k(a)$ complement of it. Let $\{x\}$ be a k basis of this complement. Then:

$$a^{-1}xa = qx \quad \text{or equivalently} \quad xa = qax \quad \text{for some } q \in k. \quad (3)$$

Since $a^N = 1$ it follows that $q^N = 1$. Set $n = |q|$. Then $n|N$. Also set $r = |q^\nu|$. By (3) $(x \otimes a^\nu)(1 \otimes x) = q^{-\nu}(1 \otimes x)(x \otimes a^\nu)$, hence by Proposition 1.2

$$\Delta(x^r) = (x \otimes a^\nu + 1 \otimes x)^r = x^r \otimes a^{\nu r} + 1 \otimes x^r.$$

Moreover,

$$x^r a^{\nu r} = q^{\nu r^2} a^{\nu r} x^r = a^{\nu r} x^r.$$

Thus it follows from Lemma 1.3 that $x^r = \alpha(a^{\nu r} - 1)$ for some $\alpha \in k$. Therefore, if either $\alpha = 0$ or $N|\nu r$ then there exists a surjection of Hopf algebras $\mathcal{H}_{n,q,N,\nu,0} \rightarrow \mathcal{H}$, which is also an injection by Proposition 2.1(3) and [M, Theorem 5.3.1], and we are done. If $\alpha \neq 0$ and $N \nmid \nu r$ then $x^r \neq 0$. Since $\alpha(a^{\nu r} - 1)$ and a commute, so do x^r and a . But, $x^r \neq 0$, hence $q^r = 1$ which is equivalent to $n|r = \frac{n}{(n,\nu)}$. Thus, $(n,\nu) = 1$ and $r = n$, and we conclude

that there exists a surjection of Hopf algebras $\mathcal{H}_{n,q,N,\nu,\alpha} \rightarrow \mathcal{H}$, which is also an injection by Proposition 2.1(3) and [M, Theorem 5.3.1]. ■

We conclude this section by recalling the definition of $U_{(N,\nu,\omega)}$ [R1, 5.2]. Let ν and N be positive integers such that $1 \leq \nu < N$ and $N \nmid \nu^2$. Suppose that $\omega \in k$ is a primitive N th root of unity. Let $q = \omega^\nu$ and $r = |q^\nu| = |\omega^{\nu^2}|$. As an algebra $U = U_{(N,\nu,\omega)}$ is generated by a, x and y which satisfy the relations

$$a^N = 1, \quad x^r = 0, \quad y^r = 0, \quad xa = qax, \quad ya = q^{-1}ay \quad \text{and} \quad yx - q^{-\nu}xy = a^{2\nu} - 1.$$

The coalgebra structure of U is determined by

$$\begin{aligned} \Delta(a) &= a \otimes a, \quad \Delta(x) = x \otimes a^\nu + 1 \otimes x, \quad \Delta(y) = y \otimes a^\nu + 1 \otimes y, \\ \varepsilon(a) &= 1, \quad \varepsilon(x) = 0 \quad \text{and} \quad \varepsilon(y) = 0. \end{aligned}$$

The antipode of U is determined by

$$s(a) = a^{-1}, \quad s(x) = -q^{-\nu}a^{-\nu}x \quad \text{and} \quad s(y) = -q^\nu a^{-\nu}y.$$

It follows from Lemma 1.1, [G2, Lemma 1.2.1] and [R1, Proposition 10 d) and Theorem 5] that U satisfies the following:

- (U.1) As an algebra U is generated by a grouplike element a of order N , and by the space $P_{a^\nu,1}(U) = sp\{a^\nu - 1, x, y\}$ of dimension 3. In particular, $G(U) = \langle a \rangle$.
- (U.2) $P_{a^\nu,1}(U) = sp_k\{a^\nu - 1\}$ for all $\nu \neq \nu$. In particular, it follows from (1) that $U_1 = sp\{a^i, a^jx, a^ly \mid 0 \leq i, j, l \leq N-1\}$.
- (U.3) U is minimal quasitriangular.
- (U.4) U is pointed, ribbon and unimodular.

The author proved that over an algebraically closed field of characteristic zero, if U is a Hopf algebra which satisfies (U.1)-(U.3) with prime N then either $U = U_{(2,1,-1)}$ or there exists a primitive N th root of unity q so that $U = U_q(sl_2)'$ [G3, Theorem 1.2.2].

2 The Family $\mathcal{H}_{n,q,N,\nu,\alpha}$

In this section we study some properties of the Hopf algebra $\mathcal{H}_{n,q,N,\nu,\alpha}$ and of its dual Hopf algebra $\mathcal{H}_{n,q,N,\nu,\alpha}^*$. Beside of being interesting on their own right they will prove to be crucial in Section 3.

Proposition 2.1 *Let $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$. Then:*

1. $\mathcal{H}_\alpha$ is a pointed Hopf algebra with $G(\mathcal{H}_\alpha) = \langle a \rangle$ a cyclic group of order N .

2. The set $\{a^i x^j | 0 \leq i \leq N-1, 0 \leq j \leq r-1\}$ forms a linear basis of $\mathcal{H}_\alpha$, hence $\dim \mathcal{H}_\alpha = Nr$.
3. If $n|\nu$ then $P_{a^\nu,1}(\mathcal{H}_0) = \text{sp}_k\{a^\nu - 1\}$. If $n \nmid \nu$ then $P_{a^\nu,1}(\mathcal{H}_\alpha) = \text{sp}_k\{a^\nu - 1, x\}$, $P_{a^i,1}(\mathcal{H}_\alpha) = \text{sp}_k\{a^i - 1\}$ for all $i \neq \nu$, and $(\mathcal{H}_\alpha)_1 = \text{sp}\{a^i, a^j x | 0 \leq i, j \leq N-1\}$.
4. The elements $\lambda_l = \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i\right) x^{r-1}$ and $\lambda_r = x^{r-1} \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i\right)$ are non-zero left and right integral of $\mathcal{H}_\alpha$. In particular $\mathcal{H}_\alpha$ is not unimodular when $r > 1$, and the distinguished grouplike element of $\mathcal{H}_\alpha^*$, ψ , is determined by $\psi(a) = q^{r-1}$ and $\psi(x) = 0$.
5. As an algebra $\mathcal{H}_\alpha \cong k\mathbb{Z}_{N/n} \#_\sigma H_{n,q,n,\nu}$ is a crossed product of a group algebra and a Radford Hopf algebra with the trivial action.
6. If the field k is algebraically closed then $\mathcal{H}_\alpha \cong \mathcal{H}_\beta$ as Hopf algebras for any $\alpha, \beta \in k^*$.

Proof: 1. Follows from Lemma 1.1.

2. For $\alpha = 0$ it was proved in [R1, Proposition 7 b)] using Bergman's Diamond Lemma (see e.g. [R1, page 214] for a description of this lemma). The proof for $\alpha \neq 0$ is very similar, and we merely sketch it following the line of argument of the proof of [R1, Proposition 7 b)]. Using the notation of Remark 1.4, let $T(C) = k\{A, B, D, X\}$ where $A < B < D < X$ and the substitution rules are:

$$A^N \leftarrow 1, \quad B \leftarrow A^\nu, \quad D \leftarrow 1, \quad XA \leftarrow qAX \quad \text{and} \quad X^n \leftarrow \alpha(A^{\nu n} - 1).$$

The only ambiguities to consider are overlap ambiguities, and they are all easily resolved. We just note that in the case $\alpha \neq 0$, it is necessary to have $r = n$ (so that $q^r = q^n = 1$) when resolving, for instance, the overlap ambiguity $(X^{n-1}X)A = X^{n-1}(XA)$.

3. For $\alpha = 0$ it was proved in [R1, Theorem 4 a)]. Suppose $\alpha \neq 0$ (so in particular $n \nmid \nu$), and let $z \in P_{a^i,1}(\mathcal{H}_\alpha)$. Then the result follows in a straightforward manner after expressing z in terms of the basis given in part 2, to compare the expressions of $\Delta(z)$ and $z \otimes a^i + 1 \otimes z$. The description of $(\mathcal{H}_\alpha)_1$ follows now from (1).

4. We show that λ_l is a left integral. For $\alpha = 0$ it was shown in [R1, Proposition 7 d)]. Suppose $\alpha \neq 0$. Since $\frac{1}{N} \left(\sum_{i=0}^{N-1} a^i\right)$ is a 2-sided integral of $k(a)$ it follows that $a\lambda_l = \lambda_l = \varepsilon(a)\lambda_l$. Now, let $\phi \in G(\mathcal{H}_\alpha^*)$ be determined by $\phi(a) = q$ and $\phi(x) = 0$. Using the facts that $q^n = 1$, and $\rightarrow$ is an algebra map we compute,

$$\begin{aligned} x\lambda_l &= x \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) x^{n-1} \\ &= \left(\frac{1}{N} \sum_{i=0}^{N-1} q^i a^i \right) x^n = \left(\frac{1}{N} \sum_{i=0}^{N-1} q^i a^i \right) \alpha(a^{\nu n} - 1) \\ &= \left(\phi \rightarrow \frac{1}{N} \sum_{i=0}^{N-1} a^i \right) \alpha(a^{\nu n} - 1) = \left(\phi \rightarrow \frac{1}{N} \sum_{i=0}^{N-1} a^i \right) (\phi \rightarrow (\alpha(a^{\nu n} - 1))) \\ &= \phi \rightarrow \left(\left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) (\alpha(a^{\nu n} - 1)) \right) = \phi \rightarrow 0 = 0 = \varepsilon(x)\lambda_l. \end{aligned}$$

Similarly, λ_r is a right integral of $\mathcal{H}_\alpha$. Since $\lambda_l a = q^{n-1} \lambda_l = q^{-1} \lambda_l$ and $\lambda_l x = 0$, it follows that ψ is the distinguished grouplike element of $\mathcal{H}_\alpha^*$ and that $\mathcal{H}_\alpha$ is not unimodular.

5. Since a^n is central in $\mathcal{H}_\alpha$ it follows that $K = k(a^n) \cong k\mathbb{Z}_{N/n}$ is a central Hopf subalgebra of $\mathcal{H}_\alpha$, hence is normal. Let $\pi : \mathcal{H}_\alpha \rightarrow H_{n,q,n,\nu}$ be the Hopf algebra surjection determined by $\pi(a) = a$ and $\pi(x) = x$. Then $\mathcal{H}_\alpha K^+ = \ker \pi$, hence the following is a short exact sequence of Hopf algebra maps:

$$K \xhookrightarrow{i} \mathcal{H}_\alpha \xrightarrow{\pi} H_{n,q,n,\nu}$$

where i is the inclusion map. Thus, $\mathcal{H}_\alpha \cong k\mathbb{Z}_{N/n} \#_\sigma H_{n,q,n,\nu}$. Finally, by [DT], the action is the trivial one since $k\mathbb{Z}_{N/n}$ is central.

6. Let $\mathcal{H}_\alpha = k \langle a, x \rangle$ and $\mathcal{H}_\beta = k \langle g, y \rangle$. Then it is straightforward to check that the map $f : \mathcal{H}_\alpha \rightarrow \mathcal{H}_\beta$ determined by $f(a) = g$ and $f(x) = (\alpha/\beta)^{1/n} y$ is an isomorphism of Hopf algebras. This completes the proof of the proposition. ■

We now consider the family of the dual Hopf algebras $\mathcal{H}_{n,q,N,\nu,\alpha}^*$.

Lemma 2.2 *Let $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$, and assume k contains a primitive N th root of unity ω . If $\alpha = 0$ then $|G(\mathcal{H}_0^*)| = N$, and if $\alpha \neq 0$ then $|G(\mathcal{H}_\alpha^*)| = \#\{0 \leq i \leq N-1 \mid N \text{ divides } \nu n i\}$.*

Proof: For $\alpha = 0$ and $n = 1$ the statement is clear since $\mathcal{H}_\alpha = k\mathbb{Z}_N$. Suppose $\alpha = 0$ and $n > 1$, and let $\phi \in G(\mathcal{H}_0^*)$. Since ϕ is an algebra homomorphism and $a^N = 1$, it follows that $\phi(a) = \omega^i$ for some $0 \leq i \leq N-1$, and the relation $xa = qax$ yields that $\phi(x) = 0$. Now, it is straightforward to check that the map $\phi : \mathcal{H}_0 \rightarrow k$ determined by $\phi(a) = \omega^i$ for some $0 \leq i \leq N-1$ and $\phi(x) = 0$ is indeed an algebra homomorphism. Thus, the result follows.

Suppose $\alpha \neq 0$ (so in particular $n > 1$), and let $\phi \in G(\mathcal{H}_\alpha^*)$. As above, $\phi(a) = \omega^i$ for some $0 \leq i \leq N-1$, and $\phi(x) = 0$. Now, since $x^n = \alpha(a^{\nu n} - 1)$ it follows that $0 = \phi(x)^n = \phi(x^n) = \alpha\phi(a^{\nu n} - 1) = \alpha(\omega^{\nu n i} - 1)$ which is equivalent to $N \mid \nu n i$. Therefore, $G(\mathcal{H}_\alpha^*)$ and $\{0 \leq i \leq N-1 \mid N \text{ divides } \nu n i\}$ are in one to one correspondence and the result follows. ■

Remark 2.3 Since $\{0 \leq i \leq N-1 \mid N \text{ divides } \nu n i\}$ is a sub group of $\mathbb{Z}_N$ it follows that $\#\{0 \leq i \leq N-1 \mid N \text{ divides } \nu n i\}$ divides N . Moreover, since $N/n \leq N-1$ is an integer we have that $n \leq |G(\mathcal{H}_\alpha^*)| \leq N$.

We are ready now to characterize Radford's $H_{(N,\nu,\omega)}$ via self duality.

Theorem 2.4 *Let $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$. Then $\mathcal{H}_\alpha$ is self dual if and only if $\alpha = 0$ and either $\mathcal{H}_0 = k\mathbb{Z}_N$ or $\mathcal{H}_0 = H_{(N,\nu,\omega)}$.*

Proof: The 'only if' part is [R1, Proposition 8]. Conversely, suppose $\mathcal{H}_\alpha$ is self dual, and assume on the contrary that $\alpha \neq 0$. Then $|G(\mathcal{H}_\alpha^*)| = |G(\mathcal{H}_\alpha)| = N$, hence by Lemma 2.2, $\#\{0 \leq i \leq N-1 \mid N \text{ divides } \nu n i\} = N$. In particular $N \mid \nu n$ which is a contradiction. Therefore, $\alpha = 0$ and the result follows from [R1, Proposition 8]. ■

We now determine when $\mathcal{H}_\alpha^*$ is pointed.

Theorem 2.5 *Let $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$, and assume k is algebraically closed. Then $\mathcal{H}_\alpha^*$ is pointed if and only if $\alpha = 0$. Moreover, if $\alpha \neq 0$ then $\mathcal{H}_\alpha^*$ contains an n^2 -dimensional simple subcoalgebra.*

Proof: Suppose $\alpha = 0$. If $r = 1$ then $\mathcal{H}_0 = k\mathbb{Z}_N$, hence $\mathcal{H}_0^* = k\mathbb{Z}_N$ is pointed. Suppose $r > 1$ (so $\mathcal{H}_0 \neq k\mathbb{Z}_N$). We show that $\mathcal{H}_0^*$ is pointed by proving that all its irreducible comodules are 1-dimensional, and hence that all its simple subcoalgebras are 1-dimensional which is to say that it is pointed. Equivalently, we prove that all the irreducible $\mathcal{H}_0$ -modules are 1-dimensional. Indeed, suppose V is an irreducible $\mathcal{H}_0$ -module so that $\dim V > 1$. Let $0 \neq v \in V$ so that $a \cdot v = \lambda v$, $\lambda \in k$. Since $a \cdot (x^i \cdot v) = q^i x^i \cdot (a \cdot v) = q^i \lambda (x^i \cdot v)$ it follows that either $x^i \cdot v = 0$ or $x^i \cdot v$ is an eigenvector of a , all i . Let $i \geq 0$ be the minimal integer so that $x^i \cdot v \neq 0$ but $x^{i+1} \cdot v = 0$ (such an i exists since $x^r = 0$). Then, $sp_k\{x^i \cdot v\}$ is stable under the action of a and x , hence is a non-trivial sub-module of V which is a contradiction, and the result follows.

Conversely, suppose $\alpha \neq 0$ (so in particular $n > 1$). We show that $\mathcal{H}_\alpha^*$ is not pointed by constructing irreducible $\mathcal{H}_\alpha$ -modules of dimension greater than 1. It will follow that $\mathcal{H}_\alpha^*$ contains simple subcoalgebras of dimension greater than 1, and hence is not pointed. Pick an N th primitive root of unity $\omega \in k$, and define the $\mathcal{H}_\alpha$ -module V_ω as follows: V_ω has a linear basis $\{v_0, \dots, v_{n-1}\}$ and the action, $\cdot$, is determined by

$$a \cdot v_i = q^{-i} \omega v_i \quad \text{and} \quad x \cdot v_i = \beta v_{i+1}$$

where $\beta = [\alpha(\omega^{\nu n} - 1)]^{1/n}$. Note that since $\omega \neq 1$ and N does not divide νn it follows that $\beta \neq 0$. It is straightforward to check that this definition is compatible with the defining relations of $\mathcal{H}_\alpha$. For example, $x^n \cdot v_i = \beta^n v_{i+n} = \alpha(\omega^{\nu n} - 1)v_i$ and $\alpha(a^{\nu n} - 1) \cdot v_i = \alpha((q^{-i} \omega)^{\nu n} - 1)v_i = \alpha(\omega^{\nu n} - 1)v_i$ as $q^n = 1$. We claim that V_ω is irreducible. Indeed, let $0 \neq V \subseteq V_\omega$ be a submodule. Since $a \cdot : V \rightarrow V$ is diagonalizable, $V_\omega = \bigoplus_{i=0}^{n-1} sp_k\{v_i\}$ and v_i, v_j are eigenvectors of distinct eigenvalues for $0 \leq i \neq j \leq n-1$, it follows that there exists $0 \leq i \leq n-1$ so that $v_i \in V$. But then, $v_j = x^{j-i} \cdot v_i \in V$ for all j , and hence $V = V_\omega$. Therefore, if $\mathcal{H}_\alpha^*$ is pointed then $\alpha = 0$. This concludes the proof of the theorem. ■

By [R1, Proposition 8], $H_{(N,\nu,\omega)}$ is self dual, so in particular its dual is pointed. By Theorem 2.5, $H_{n,q,N,\nu}^*$ is always pointed regardless being self dual. In fact we can say more.

Proposition 2.6 *Let $H = H_{n,q,N,\nu} \neq k\mathbb{Z}_N$, and let $\omega \in k$ be a primitive N th root of unity. Then $H^* = H_{\frac{N}{(N,\nu)},\omega^\nu,N,\mu}$ for some $0 \leq \mu \leq N-1$ so that $q = \omega^\mu$.*

Proof: Since $n|N$ there exists $0 \leq \mu \leq N-1$ so that $q = \omega^\mu$. Using Proposition 2.1 (2), define $A, X \in H^*$ as follows:

$$\langle A, a^i x^j \rangle = \delta_{j,0} \omega^i \quad \langle X, a^i x^j \rangle = \delta_{j,1}$$

for all $0 \leq i \leq N-1, 0 \leq j \leq r-1$. Then it is straightforward to check that $A \in G(H^*)$ is of order N , $X \in P_{A^\mu, \varepsilon}(H^*)$, $X^r = 0$, $XA = \omega^\nu AX$, and finally that $H^* = k \langle A, X \rangle = H_{\frac{N}{(N,\nu)},\omega^\nu,N,\mu}$. ■

Corollary 2.7 Let $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha} \neq k\mathbb{Z}_N$, and let $\omega \in k$ be a primitive N th root of unity. Then as an algebra $\mathcal{H}_\alpha^* \cong H_{\frac{n}{(n,\nu)},\omega^\nu,n,\mu} \#_\tau k\mathbb{Z}_{N/n}$ is a crossed product of a Radford Hopf algebra and a group algebra for some $0 \leq \mu \leq N-1$ so that $q = \omega^\mu$. If moreover $\alpha \neq 0$ then $\mathcal{H}_\alpha^* \cong H_{n,q,n,\nu} \#_\tau k\mathbb{Z}_{N/n}$.

Proof: By the proof of Proposition 2.1 (5), we have the following short exact sequence of Hopf algebra maps:

$$k\mathbb{Z}_{N/n} \xrightarrow{i} \mathcal{H}_\alpha \xrightarrow{\pi} H_{n,q,n,\nu}.$$

Dualizing yields the following short exact sequence of Hopf algebra maps:

$$H_{n,q,n,\nu}^* \xrightarrow{\pi^*} \mathcal{H}_\alpha^* \xrightarrow{i^*} (k\mathbb{Z}_{N/n})^*.$$

Since $(k\mathbb{Z}_{N/n})^* \cong k\mathbb{Z}_{N/n}$ as Hopf algebras and by Proposition 2.6, $H_{n,q,n,\nu}^* \cong H_{\frac{n}{(n,\nu)},\omega^\nu,n,\mu}$ for some $0 \leq \mu \leq N-1$ so that $q = \omega^\mu$, it follows that $\mathcal{H}_\alpha^* \cong H_{\frac{n}{(n,\nu)},\omega^\nu,n,\mu} \#_\tau k\mathbb{Z}_{N/n}$. If moreover $\alpha \neq 0$ then $(n,\nu) = 1$ and hence $\omega = q^{\nu^{-1}(\text{mod } n)}$ is a primitive n th root of unity so that $\omega^\nu = q$. Thus, by Theorem 2.4, $H_{n,q,n,\nu}^* \cong H_{n,q,n,\nu}$ and the result follows. ■

We shall need the following lemma in which we find all the Hopf subalgebras of $H_{n,q,N,\nu}$.

Lemma 2.8 Let $H = H_{n,q,N,\nu} \neq k\mathbb{Z}_N$ and $B \subseteq H$ be a Hopf subalgebra of H . Then either $B = kG(B)$ where $G(B)$ is a subgroup of (a) , or there exists an integer $1 \leq s \leq N$ so that $s|N$, $(N/s)|\nu$ and $B = k \langle a^{N/s}, x \rangle = H_{\frac{n}{(n,N/s)},q^{N/s},s,\frac{\nu}{N/s}}$.

Proof: Suppose $B \neq kG(B)$. Then by Taft-Wilson Theorem [TW], B must contain a non-trivial skew primitive element P . Without loss of generality we may assume that $P \in P_{a^i,1}(B)$ for some $0 \leq i \leq N-1$. By Proposition 2.1(3), $i = \nu$, and hence $x \in B$.

Let $i : B \rightarrow H$ be the inclusion map. Then $i^* : H^* \rightarrow B^*$ is a surjection of Hopf algebras. By Proposition 2.6, $H^* = k \langle A, X \rangle = H_{\frac{N}{(N,\nu)},\omega^\nu,N,\mu}$ for some $0 \leq \mu \leq N-1$ so that $q = \omega^\mu$ (where A, X are as in the proof of Proposition 2.6). Therefore, B^* is generated as an algebra by $i^*(A)$, $i^*(X)$ and $i^*(A) \in G(B^*)$, $i^*(X) \in P_{i^*(A)^\mu,1}(B^*)$ (note that since $x \in B$, $i^*(X)$ must be non-trivial). Set $s = |i^*(A)|$. Then $s|N$. It is now straightforward to check that there exists a surjection of Hopf algebras $H_{\frac{N}{(N,\nu)},\omega^\nu,s,\mu} \rightarrow B^* = k \langle i^*(A), i^*(X) \rangle$ (here μ means $\mu(\text{mod } s)$) which is also an injection by Proposition 2.1(3) and [M, Theorem 5.3.1]. Now, $\omega^{N/s}$ is a primitive s th root of unity, and hence by Proposition 2.6, $B = H_{\frac{N}{(N,\nu)},\omega^\nu,s,\mu}^* = H_{\frac{s}{(s,\mu)},q^{N/s},s,\beta}$ for some β so that $\omega^\nu = \omega^{(N\beta)/s}$. In particular the last equation implies that $(N/s)|\nu$ and $\beta = \frac{\nu}{N/s}(\text{mod } s)$. This concludes the proof of the lemma. ■

It is known that $H_{(N,\nu,\omega)}$ is quasitriangular if and only if $N = 2\nu$ and ν is odd [G1,R1]. In the following we investigate the quasitriangularity of $\mathcal{H}_\alpha$.

Theorem 2.9 Let k be an algebraically closed field of characteristic 0. Then $\mathcal{H}_\alpha = \mathcal{H}_{n,q,N,\nu,\alpha}$ is quasitriangular if and only if either $\mathcal{H}_\alpha = k\mathbb{Z}_N$ or $\mathcal{H}_\alpha = H_{(2\nu,\nu,\omega)}$, where ν is odd.

Proof: We have to prove the 'only if' part. We first show that if $\mathcal{H}_\alpha$ is quasitriangular then $\alpha = 0$. Suppose on the contrary that $(\mathcal{H}_\alpha, R)$ is quasitriangular and $\alpha \neq 0$. Write $R = \sum R^{(1)} \otimes R^{(2)}$ in the shortest possible way. Then $A = sp\{R^{(1)}\}$ and $B = sp\{R^{(2)}\}$ are Hopf subalgebras of $\mathcal{H}_\alpha$ and $A \cong B^{*cop}$. In particular A, B, A^* and B^* are pointed. Suppose $A \neq kG(A)$. Then using similar arguments to those used in the first paragraph of the proof of Lemma 2.8 yields that $a^\nu, x \in A$. Therefore, we have an inclusion of Hopf algebras $k \langle a^\nu, x \rangle \subseteq A$, and hence a surjection of Hopf algebras $A^* \rightarrow k \langle a^\nu, x \rangle^*$. Since A^* is pointed so is $k \langle a^\nu, x \rangle^*$. But, $k \langle a^\nu, x \rangle^* = \mathcal{H}_{n, q^\nu, \frac{N}{(N, \nu)}, 1, \alpha}$ is *not* pointed by Theorem 2.5, which is a contradiction. We conclude that $A = kG(A) = k(a^m)$ for some $m|N$, and since $B \cong A^{*cop}$ it follows that $B = k(a^m)$ as well. Therefore, $R \in k(a) \otimes k(a)$. Let $\omega \in k$ be a primitive N th root of unity. By [R1, page 219], there exists $0 \leq l \leq N-1$ so that

$$R = \frac{1}{N} \left(\sum_{i,j=0}^{N-1} \omega^{-ij} a^i \otimes a^{jl} \right). \quad (4)$$

By (QT.5), $(a^\nu \otimes x + x \otimes 1)R = R(x \otimes a^\nu + 1 \otimes x)$; that is,

$$\begin{aligned} \frac{1}{N} \left(\sum_{i,j=0}^{N-1} \omega^{-ij} q^{jl} a^{i+\nu} \otimes a^{jl} x \right) + \frac{1}{N} \left(\sum_{i,j=0}^{N-1} \omega^{-ij} q^i a^i x \otimes a^{jl} \right) = \\ \frac{1}{N} \left(\sum_{i,j=0}^{N-1} \omega^{-ij} a^i x \otimes a^{jl+\nu} \right) + \frac{1}{N} \left(\sum_{i,j=0}^{N-1} \omega^{-ij} a^i \otimes a^{jl} x \right). \end{aligned} \quad (5)$$

Thus, $\frac{1}{N} \omega^{-ij} q^{jl} = \frac{1}{N} \omega^{-(i+\nu)j}$, i.e. $q^{jl} = \omega^{-\nu j}$ for all j . In particular, for $j = n$ we obtain $1 = q^{nl} = \omega^{-\nu n}$. Thus, $N|\nu n$ which is a contradiction. Therefore $\alpha = 0$.

Suppose $H_{n,q,N,\nu} \neq k\mathbb{Z}_N$, $(H_{n,q,N,\nu}, R)$ is quasitriangular, and let A and B be defined as before. Suppose $A \neq kG(A)$ (equivalently, $B \neq kG(B)$ since $A \cong B^{*cop}$). By Lemma 2.8, there exist integers $1 \leq s, s' \leq N$ so that $s, s'|N$, $(N/s), (N/s')|\nu$ and

$$A = k \langle a^{N/s}, x \rangle = H_{\frac{n}{(n, N/s)}, q^{N/s}, s, \frac{\nu}{N/s}}$$

and

$$B = k \langle a^{N/s'}, y \rangle = H_{\frac{n}{(n, N/s')}, q^{N/s'}, s', \frac{\nu}{N/s'}}.$$

Since $A \cong B^{*cop}$, it follows from Proposition 2.6 that $s = s'$, and hence by [G2, Lemma 1.1.2], there exists a primitive sth root of unity χ so that $q^{N/s} = \chi^{\frac{\nu}{N/s}}$ and $q^{N/s} = \chi^{-\frac{\nu}{N/s}}$. Therefore $N|2\nu$, and since $\nu < N$, we conclude that $N = 2\nu$. Now, since A and B generate $H_{n,q,N,\nu}$ as a Hopf algebra, we must have $s = N$. Therefore we may assume that $\chi = \omega$ and hence $q = \omega^\nu$. Since by assumption $H_{n,q,N,\nu} \neq k\mathbb{Z}_N$, we have that $1 < r = |q^\nu| = |\omega^{\nu^2}|$, and hence $2\nu \nmid \nu^2$ which implies that ν is odd. Therefore, $A = B = H_{n,q,N,\nu} = H_{(2\nu, \nu, \omega)}$ where ν is odd.

If $A = B = k(a^m)$ then again R must be of the form (4). But then it follows from (5) that l must be invertible in $\mathbb{Z}_N$, and hence that on one hand $q = \omega^{-\nu l^{-1}}$ and on the other hand, $q = \omega^{\nu l^{-1}}$. Therefore, $N = 2\nu$ and $q = \omega^\nu = -1$. By [R1, Corollary 3(a)], $H_{n,q,N,\nu} = H_{(2\nu, \nu, \omega)}$ where ν is odd. This completes the proof of the theorem. ■

We conclude this section with an example.

Example 2.10 The least dimensional new Hopf algebra in our family is an 8–dimensional Hopf algebra, namely $H_8 = \mathcal{H}_{2,-1,4,1,1}$, generated as an algebra by $a \in G(H_8)$ and $x \in P_{a,1}(H_8)$ which satisfy the relations

$$a^4 = 1, \quad x^2 = a^2 - 1 \quad \text{and} \quad xa = -ax.$$

By Proposition 2.1, as an algebra $H_8 = k\mathbb{Z}_2 \#_{\sigma} H_4$ is an extension of $k\mathbb{Z}_2$ by Sweedler’s Hopf algebra H_4 . By Theorem 2.9, H_8 is not quasitriangular even though $k\mathbb{Z}_2$ and H_4 are. By Lemma 2.2, $G(H_8^*) = \{1, \phi\}$ is of order 2, where $\phi : H_8 \rightarrow k$ is determined by $\phi(a) = -1$ and $\phi(x) = 0$. By Corollary 2.7, as an algebra $H_8^* = H_4 \#_{\tau} k\mathbb{Z}_2$ is an extension of H_4 by $k\mathbb{Z}_2$. By Theorem 2.5, H_8^* is not pointed and it contains a (unique) 4–dimensional simple sub coalgebra. In particular H_8 is not self dual.

3 The Family $\mathcal{U}_{(n,N,\nu,q,\alpha,\beta,\gamma)}$

In this section we first construct a new family of finite dimensional pointed and unimodular Hopf algebras, denoted by $\mathcal{U}_{(n,N,\nu,q,\alpha,\beta,\gamma)}$, which generalizes Radford’s family $U_{(N,\nu,\omega)}$. Second, we show that over any infinite field which contains a primitive n th root of unity, our new family contains infinitely many non-isomorphic Hopf algebras of any dimension of the form Nn^2 , where $2 < n < N$ are integers so that n divides N . Thus, we prove that Kaplansky’s 10th conjecture [K] is false. Third, we use Theorem 2.5 to generalize [G3, Theorem 1.2.2] and show that over an algebraically closed field of characteristic zero the family $U_{(N,\nu,\omega)}$, any N , is unique with respect to properties (U.1)-(U.3).

We start by introducing $\mathcal{U}_{(n,N,\nu,q,\alpha,\beta,\gamma)}$. Let $n > 1$, ν and N be positive integers such that $1 \leq \nu < N$ and $n|N$, and let $\alpha, \beta, \gamma \in k$. Suppose that $q \in k$ is a primitive n th root of unity, and let $r = |q^{\nu}|$. For $\alpha = \beta = 0$ we define $\mathcal{U}_{(n,N,\nu,q,0,0,\gamma)}$ exactly as $U_{(N,\nu,\omega)}$ except that now $yx - q^{-\nu}xy = \gamma(a^{2\nu} - 1)$. For $(\alpha, \beta) \neq (0, 0)$ we assume that $(n, \nu) = 1$ (hence $r = n$), and $N \nmid \nu n$. In this case, as an algebra $\mathcal{U}_{\alpha,\beta,\gamma} = \mathcal{U}_{(n,N,\nu,q,\alpha,\beta,\gamma)}$ is generated by a, x and y which satisfy the relations

$$\begin{aligned} a^N &= 1, \quad x^n = \alpha(a^{\nu n} - 1), \quad y^n = \beta(a^{\nu n} - 1), \\ xa &= qax, \quad ya = q^{-1}ay \quad \text{and} \quad yx - q^{-\nu}xy = \gamma(a^{2\nu} - 1). \end{aligned}$$

The coalgebra structure of $\mathcal{U}_{\alpha,\beta,\gamma}$ is determined by

$$\begin{aligned} \Delta(a) &= a \otimes a, \quad \Delta(x) = x \otimes a^{\nu} + 1 \otimes x, \quad \Delta(y) = y \otimes a^{\nu} + 1 \otimes y, \\ \varepsilon(a) &= 1, \quad \varepsilon(x) = 0 \quad \text{and} \quad \varepsilon(y) = 0. \end{aligned}$$

The antipode of $\mathcal{U}_{\alpha,\beta,\gamma}$ is determined by

$$s(a) = a^{-1}, \quad s(x) = -q^{-\nu}a^{-\nu}x \quad \text{and} \quad s(y) = -q^{\nu}a^{-\nu}y.$$

Note that if ω is a primitive N th root of unity and $N \nmid \nu^2$, then $U_{(N,\nu,\omega)} = \mathcal{U}_{(\frac{N}{(N,\nu)}, N, \nu, \omega^{\nu}, 0, 0, 1)}$, and $\mathcal{U}_{(\frac{N}{(N,\nu)}, N, \nu, \omega^{\nu}, 0, 0, \gamma)} \cong U_{(N,\nu,\omega)}$ as Hopf algebras for any $\gamma \in k^*$ which has a square root in k .

Remark 3.1 Suppose $(\alpha, \beta) \neq (0, 0)$. Then to construct $\mathcal{U}_{(n, N, \nu, q, \alpha, \beta, \gamma)}$ we follow the line of argument and notations used by Radford in order to construct $U_{(N, \nu, \omega)}$ [R1, 5.2] (see also Remark 1.4). We let C be the 5-dimensional coalgebra over k with basis $\{A, B, D, X, Y\}$ whose structure is determined by

$$\Delta(A) = A \otimes A, \quad \Delta(B) = B \otimes B, \quad \Delta(D) = D \otimes D,$$

$$\Delta(X) = X \otimes B + D \otimes X \quad \text{and} \quad \Delta(Y) = Y \otimes B + D \otimes Y,$$

and let I be the ideal of the tensor algebra $T(C)$ generated by

$$A^N - 1, \quad B - A^\nu, \quad D - 1, \quad XA - qAX, \quad YA - q^{-1}AY,$$

$$X^n - \alpha(A^{\nu n} - 1), \quad Y^n - \beta(A^{\nu n} - 1) \quad \text{and} \quad YX - q^{-\nu}XY - \gamma(A^{2\nu} - 1).$$

The details are straightforward to work out.

Proposition 3.2 *Let $\mathcal{U}_{\alpha, \beta, \gamma} = \mathcal{U}_{(n, N, \nu, q, \alpha, \beta, \gamma)}$. Then:*

1. $\mathcal{U}_{\alpha, \beta, \gamma}$ is a pointed Hopf algebra with $G(\mathcal{U}_{\alpha, \beta, \gamma}) = (a)$ a cyclic group of order N .
2. The set $\{a^i x^j y^l \mid 0 \leq i \leq N-1, 0 \leq j, l \leq r-1\}$ forms a linear basis of $\mathcal{U}_{\alpha, \beta, \gamma}$, hence $\dim \mathcal{U}_{\alpha, \beta, \gamma} = Nr^2$.
3. $P_{a^\nu, 1}(\mathcal{U}_{\alpha, \beta, \gamma}) = \text{sp}_k\{a^\nu - 1, x, y\}$, $P_{a^v, 1}(\mathcal{U}_{\alpha, \beta, \gamma}) = \text{sp}_k\{a^v - 1\}$ for all $v \neq \nu$, and $(\mathcal{U}_{\alpha, \beta, \gamma})_1 = \text{sp}_k\{a^i, a^j x, a^l y \mid 0 \leq i, j, l \leq N-1\}$.
4. The element $\lambda = \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i\right) x^{r-1} y^{r-1}$ is a non-zero 2-sided integral of $\mathcal{U}_{\alpha, \beta, \gamma}$. In particular $\mathcal{U}_{\alpha, \beta, \gamma}$ is unimodular.
5. If $\alpha, \beta, \gamma \neq 0$ and $\sqrt{\gamma} \in k$, then as an algebra, $\mathcal{U}_{\alpha, \beta, \gamma} \cong k\mathbb{Z}_{N/n} \#_{\sigma} U_{(n, \nu, \omega)}$ is a crossed product of a group algebra and a Radford Hopf algebra for some primitive n th root of unity ω .
6. Let $\alpha, \beta, \gamma, \alpha', \beta', \gamma' \in k^*$. Then:

(a) $\mathcal{U}_{\alpha, \beta, 0} \cong \mathcal{U}_{\alpha', \beta', 0}$ as Hopf algebras.

(b) Suppose $N \nmid 2\nu$. Then $\mathcal{U}_{\alpha, \beta, \gamma} \cong \mathcal{U}_{\alpha', \beta', \gamma'}$ as Hopf algebras if and only if $\frac{\alpha\beta}{\gamma^n} = \frac{\alpha'\beta'}{\gamma'^n}$.

Proof: 1. Follows from Lemma 1.1.

2. For $U_{(N, \nu, \omega)}$ it was proved in [R1, Proposition 10 b)] using Bergman's Diamond Lemma. The proof for $\mathcal{U}_{\alpha, \beta, \gamma}$ is very similar, and we merely sketch it following the line of argument from the proof of [R1, Proposition 10 b)]. Using the notation of Remark 3.1, let $T(C) = k\langle A, B, D, X, Y \rangle$ where $A < B < D < X < Y$ and the substitution rules are:

$$A^N \leftarrow 1, \quad B \leftarrow A^\nu, \quad D \leftarrow 1, \quad XA \leftarrow qAX, \quad YA \leftarrow q^{-1}AY,$$

$$X^n \leftarrow \alpha(A^{\nu n} - 1), \quad Y^n \leftarrow \beta(A^{\nu n} - 1) \quad \text{and} \quad YX \leftarrow q^{-\nu}XY + \gamma(A^{2\nu} - 1).$$

The only ambiguities to consider are overlap ambiguities, and they are all easily resolved. We just note that in the case $(\alpha, \beta) \neq (0, 0)$ it is necessary to have $r = n$ (so that $q^r = q^n = 1$) when resolving, for instance, the overlap ambiguity $(YX)X^{n-1} = Y(XX^{n-1})$.

3. The proof follows the line of argument and notations of the proof of [G2, Lemma 1.2.1]. Specifically, let $A = k(a)$, $B = k < a, x >$, $H = k < a, y >$, $\{B_m\}$ be the coradical filtration of B and $\{H_m\}$ be the coradical filtration of H . Note that $B_0 = H_0 = A$. Set

$$\begin{aligned} B(1) &= Ax, & H(1) &= Ay, \\ B(m) &= B(1)^m, & H(m) &= H(1)^m \quad \text{for } 1 \leq m \\ \text{and } B(0) &= H(0) = A. \end{aligned}$$

Using parts 2 and 3 of Proposition 2.1 it is straightforward to verify that:

$$\begin{aligned} B &= \bigoplus_{i=0}^{r-1} B(i), & B(0) &= B_0, & B_1 &= B(0) \oplus B(1), \\ H &= \bigoplus_{i=0}^{r-1} H(i), & H(0) &= H_0 \quad \text{and} \quad H_1 &= H(0) \oplus H(1). \end{aligned}$$

Therefore, B and H are coradically graded [CM, Section 2]. Since $\mathcal{U}_{\alpha, \beta, \gamma} = B \otimes_A H$ it follows from [CM, Lemma 2.3] that $\mathcal{U}_{\alpha, \beta, \gamma}$ is coradically graded where,

$$\mathcal{U}_{\alpha, \beta, \gamma}(m) = \sum_{i=0}^m B(i) \otimes_A H(m-i).$$

In particular, $(\mathcal{U}_{\alpha, \beta, \gamma})_0 = \mathcal{U}_{\alpha, \beta, \gamma}(0) = A$ and

$$\mathcal{U}_{\alpha, \beta, \gamma}(1) = B(0) \otimes_A H(1) + B(1) \otimes_A H(0) = H(1) + B(1).$$

Thus $(\mathcal{U}_{\alpha, \beta, \gamma})_1 = \mathcal{U}_{\alpha, \beta, \gamma}(0) \oplus \mathcal{U}_{\alpha, \beta, \gamma}(1) = A \oplus Ax \oplus Ay$ and the result follows.

4. Since $\frac{1}{N} \left(\sum_{i=0}^{N-1} a^i \right)$ is a 2-sided integral of $k(a)$, and a commutes with $x^{r-1}y^{r-1}$ it follows that $a\lambda = \lambda a = \lambda = \varepsilon(a)\lambda$. By the same reason, since at any rate $x^r, y^r \in k(a) \cap \ker(\varepsilon)$ it follows that $x\lambda = \lambda y = 0$ (see the proof of Proposition 2.1 (4)). We show now that $\lambda x = 0$. Indeed, by simple induction we have

$$y^i x = q^{-i\nu} x y^i + \gamma \left(q^{-(i-1)\nu} \sum_{j=0}^{i-1} q^{-j\nu} \right) a^{2\nu} y^{i-1} - \gamma \left(\sum_{j=0}^{i-1} q^{-j\nu} \right) y^{i-1}.$$

Using this, and the fact that $x^r \in k(a) \cap \ker(\varepsilon)$ we compute:

$$\begin{aligned} \lambda x &= \left(\left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) x^{r-1} y^{r-1} \right) x \\ &= \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) x^{r-1} \left(q^\nu x y^{r-1} - \gamma q^{3\nu} a^{2\nu} y^{r-2} + \gamma q^\nu y^{r-2} \right) \end{aligned}$$

$$\begin{aligned}
&= \gamma \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) x^{r-1} (-q^{3\nu} a^{2\nu} + q^\nu 1) y^{r-2} \\
&= \gamma \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) (-q^\nu a^{2\nu} + q^\nu 1) x^{r-1} y^{r-2} \\
&= \varepsilon (-q^\nu a^{2\nu} + q^\nu 1) \gamma \left(\frac{1}{N} \sum_{i=0}^{N-1} a^i \right) x^{r-1} y^{r-2} = 0 = \varepsilon(x)\lambda.
\end{aligned}$$

Similarly, $y\lambda = 0$, and the result follows.

5. Since a^n is central in $\mathcal{U}_{\alpha,\beta,\gamma}$ it follows that $K = k(a^n) \cong k\mathbb{Z}_{N/n}$ is a central Hopf subalgebra of $\mathcal{U}_{\alpha,\beta,\gamma}$, hence is normal. Since $(n, \nu) = 1$ there exist integers a, b such that $1 = an + b\nu$. Set $\omega = q^b$. Then ω is a primitive n th root of unity, and $\omega^\nu = q$. Let $\pi : \mathcal{U}_{\alpha,\beta,\gamma} \rightarrow U_{(n,\nu,\omega)}$ be the Hopf algebra surjection determined by $\pi(a) = a$, $\pi(x) = \sqrt[n]{\gamma}x$ and $\pi(y) = \sqrt[n]{\gamma}y$. Then $\mathcal{U}_{\alpha,\beta,\gamma}K^+ = \ker \pi$, hence the following is a short exact sequence of Hopf algebra maps:

$$K \xrightarrow{i} \mathcal{U}_{\alpha,\beta,\gamma} \xrightarrow{\pi} U_{(n,\nu,\omega)}$$

where i is the inclusion map, and the result follows.

6. First note that since the ground field contains a primitive N th root of unity and n divides N , it also contains a primitive n th root of unity. Now, following the lines of the proof of Proposition 2.1 (6), it is straightforward to check, using part 3 of this proposition, that if $f : \mathcal{U}_{\alpha,\beta,\gamma} = k \langle a, x, y \rangle \rightarrow \mathcal{U}_{\alpha',\beta',\gamma'} = k \langle g, z, w \rangle$ is an isomorphism of Hopf algebras then $f(a) = g$, $f(x) = u(\frac{\alpha}{\alpha'})^{1/n}z$ and $f(y) = v(\frac{\beta}{\beta'})^{1/n}w$ for some n th roots of unity $u, v \in k$. If $\gamma = \gamma' = 0$ then such an f is indeed an isomorphism of Hopf algebras. If $\gamma, \gamma' \neq 0$ then since $N/2\nu$ it follows that $g^{2\nu} - 1 \neq 0$, and hence that $f(yx - q^{-\nu}xy) = \gamma f(a^{2\nu} - 1)$ if and only if $uv(\frac{\alpha\beta}{\alpha'\beta'})^{1/n}(zw - q^{-\nu}wz) = \gamma(g^{2\nu} - 1)$ if and only if $u^{-1}v^{-1}(\frac{\alpha\beta}{\alpha'\beta'})^{-1/n}\gamma = \gamma'$ if and only if $\frac{\alpha\beta}{\gamma^n} = \frac{\alpha'\beta'}{\gamma'^n}$. ■

Corollary 3.3 *Kaplansky's 10th conjecture is false over any infinite field which contains a primitive n th root of unity for some $n > 2$.*

Proof: The conjecture was that there exist only finitely many non-isomorphic Hopf algebras of any given dimension. But, since the field is infinite, it follows by Proposition 3.2 (6) that the family $\mathcal{U}_{\alpha,\beta,\gamma}$, where $\alpha, \beta, \gamma \neq 0$, contains infinitely many non-isomorphic Hopf algebras of any dimension of the form Nn^2 , where $2 < n < N$ are integers so that n divides N . ■

In what follows we characterize Radford's $U_{(N,\nu,\omega)}$ as a distinguished sub-family of our new family via minimal quasitriangularity.

Lemma 3.4 *Let k be an algebraically closed field of characteristic zero. Let U be a Hopf algebra over k which satisfies the following:*

1. $G(U) = \langle a \rangle$ is a cyclic group of order N .
2. There exists an integer $1 \leq \nu < N$ so that $\dim_k P_{a^\nu, 1}(U) = 3$.

Then U contains two Hopf subalgebras, $\mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1} = k \langle a, x \rangle$ and $\mathcal{H}_{n_2, \omega_2, N, \nu, \alpha_2} = k \langle a, y \rangle$ for some $n_1, n_2 \in \mathbb{N}$, $\alpha_1, \alpha_2, \omega_1, \omega_2 \in k$. Moreover, $\mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1} \cap \mathcal{H}_{n_2, \omega_2, N, \nu, \alpha_2} = kG(U)$.

Proof: Since $kG(U)$ is a commutative semisimple algebra over an algebraically closed field of characteristic zero all $kG(U)$ -modules are direct sums of 1-dimensional $kG(U)$ -submodules. Since $kG(U)$ acts on $P_{a^\nu, 1}(U)$ by conjugation it follows that $P_{a^\nu, 1}(U)$ is a direct sum of three 1-dimensional $kG(U)$ -submodules. Therefore we may assume that there exists a basis $\{a^\nu - 1, x, y\}$ for $P_{a^\nu, 1}(U)$ so that

$$xa = \omega_1 ax \text{ and } ya = \omega_2 ay$$

where ω_i is an N th root of unity for $i = 1, 2$. Set $n_i = |\omega_i|$ and $r_i = |\omega_i^\nu|$ for $i = 1, 2$. Following the line of argument of the proof of Theorem 1.5 yields that $x^{r_1} = \alpha_1(a^{\nu r_1} - 1)$ and $y^{r_2} = \alpha_2(a^{\nu r_2} - 1)$ for some $\alpha_1, \alpha_2 \in k$. Therefore there exist surjections of Hopf algebras $\mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1} \rightarrow k \langle a, x \rangle$ and $\mathcal{H}_{n_2, \omega_2, N, \nu, \alpha_2} \rightarrow k \langle a, y \rangle$ which are also injections by Proposition 2.1(3) and [M, Theorem 5.3.1]. This completes the proof of the lemma. ■

Proposition 3.5 *Let U be a finite dimensional Hopf algebra over an algebraically closed field k of characteristic zero. Then the following are equivalent:*

1. U satisfies the following conditions:

- (i) *There exist integers $1 \leq \nu < N$, $N \nmid \nu^2$, so that as an algebra U is generated by a grouplike element a , of order N , and by the space $P_{a^\nu, 1}(U)$ of dimension 3.*
- (ii) *$U \cong U^{* \text{ cop}}$ as Hopf algebras.*

2. $U = U_{(2\nu, \nu, \omega)}$, where ν is odd.

Proof: Suppose 2 holds. Then (i) is just (U.1). We prove (ii). By Proposition 3.2(2), the set $\{a^i x^j y^l \mid 0 \leq i < N, 0 \leq j, l < r\}$ forms a linear basis of U . Define A, X and Y in U^* as follows:

$$\langle A, a^i x^j y^l \rangle = \omega^i \delta_{j,0} \delta_{l,0}, \quad \langle X, a^i x^j y^l \rangle = \omega^{\nu i} \delta_{j,1} \delta_{l,0} = (-1)^i \delta_{j,1} \delta_{l,0}$$

and

$$\langle Y, a^i x^j y^l \rangle = \omega^{\nu i} \delta_{j,0} \delta_{l,1} = (-1)^i \delta_{j,0} \delta_{l,1}$$

for $0 \leq i \leq 2\nu - 1$ and $0 \leq j, l \leq r - 1$. Using a similar argument to the one used in the proof of [G2, Lemma 1.1.2] yields that the set $\{A^i X^j Y^l \mid 0 \leq i < N, 0 \leq j, l < r\}$ forms a linear basis of U^* . It is now straightforward to check that the map $f : U \rightarrow U^{* \text{ cop}}$ given by

$$f(a) = A, \quad f(x) = X \text{ and } f(y) = Y$$

is an isomorphism of Hopf algebras.

Assume 1 holds. It follows from Lemma 3.4 that there exist $x, y \in P_{a^\nu, 1}(U)$ such that $k \langle a, x \rangle = \mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1}$ and $k \langle a, y \rangle = \mathcal{H}_{n_2, \omega_2, N, \nu, \alpha_2}$. Let $f : U \rightarrow U^{* \text{ cop}}$ be the isomorphism of Hopf algebras which exists by (ii). Set $f(a) = A$, $f(x) = X$ and $f(y) = Y$. It follows that

$A \in G(U^*)$ and $\omega = \langle A, a \rangle \in k$ is a primitive N th root of unity. By Lemma 3.4 there exists an inclusion of Hopf algebras, $i : \mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1} \rightarrow U$, hence there exists a surjection of Hopf algebras, $i^* : U^* \rightarrow (\mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1})^*$. Since $i^*(A)$, $i^*(X)$ and $i^*(Y)$ generate $(\mathcal{H}_{n_1, \omega_1, N, \nu, \alpha_1})^*$ as an algebra it follows that either $i^*(X) \neq 0$ or $i^*(Y) \neq 0$. Therefore using a similar argument to the one used in the third paragraph of the proof of [G2, Lemma 1.1.2] we conclude that either $\langle X, x \rangle \neq 0$ or $\langle Y, x \rangle \neq 0$. Let Z be X or Y so that $\langle Z, x \rangle \neq 0$. Then $\omega_1 \langle Z, ax \rangle = \langle Z, xa \rangle$ yields $\omega_1 = \omega^{-\nu}$. In the same manner either $\langle X, y \rangle \neq 0$ or $\langle Y, y \rangle \neq 0$ and the equality $ya = \omega_2 ay$ yields $\omega_2 = \omega^{-\nu}$. In particular $\omega_1 = \omega_2$. Therefore $\langle ZA, x \rangle = \langle \omega^{-\nu} AZ, x \rangle$ implies $\omega^{-\nu} = \omega^\nu$. Thus, $\omega^{2\nu} = 1$ and $N \mid 2\nu$. Since $\nu < N$ we have $N = 2\nu$. But $N \nmid \nu^2$ hence we conclude that ν is odd. Moreover, since $\omega^\nu = -1$ it follows from the definition of $\Delta(x)$ and $\Delta(y)$ that $\Delta(yx + xy) = (xy + yx) \otimes 1 + 1 \otimes (xy + yx)$. By Lemma 1.3, $xy + yx = 0$. Finally, $r = |\omega^{\nu^2}| = \frac{2\nu}{(2\nu, \nu^2)} = 2$, hence $x^2 = \alpha_1(a^{2\nu} - 1) = 0$. Similarly, $y^2 = 0$. We conclude that there exists a surjection of Hopf algebras $U_{(2\nu, \nu, \omega)} \rightarrow U$, which is also an injection by (U.2) and [M, Theorem 5.3.1]. This completes the proof of the proposition. ■

We are ready now to characterize Radford's $U_{(N, \nu, \omega)}$ via minimal quasitriangularity.

Theorem 3.6 *Let k be an algebraically closed field of characteristic zero. Let U be a Hopf algebra which satisfies the following:*

1. *There exist integers $1 \leq \nu < N$, $N \nmid \nu^2$, so that as an algebra U is generated by a grouplike element a , of order N , and by the space $P_{a^\nu, 1}(U)$ of dimension 3.*
2. *For any positive integer $i \neq \nu \pmod{N}$, $\dim_k P_{a^i, 1}(U) = 1$.*
3. *U is minimal quasitriangular.*

Then there exists a primitive N th root of unity $\omega \in k$ such that $U = U_{(N, \nu, \omega)}$.

Proof: By Lemma 3.4, there exist $x, y \in P_{a^\nu, 1}(U)$ so that the set $\{a^\nu - 1, x, y\}$ forms a linear basis for $P_{a^\nu, 1}(U)$ and the following relations are satisfied:

$$a^N = 1, \quad xa = \omega_2 ax, \quad ya = \omega_1 ay, \quad x^{r_2} = \alpha_2(a^{\nu r_2} - 1) \text{ and } y^{r_1} = \alpha_1(a^{\nu r_1} - 1)$$

where ω_i is a primitive n_i th root of unity and $r_i = |\omega_i^\nu|$ for $i = 1, 2$. Moreover, as an algebra U is generated by a, x and y . Since U is minimal quasitriangular it follows by [R2, Proposition 2(d) and Theorem 1] that there exist two Hopf subalgebras B and H of U such that U is generated by B and H and $B^{* \text{ cop}} \cong H$ as Hopf algebras. In particular $B \neq kG(B)$ and $H \neq kG(H)$, otherwise $B = H = U \subseteq k(a)$ which is a contradiction. Note also that B^* and H^* are necessarily pointed Hopf algebras.

Let $C \subseteq U$ be a Hopf subalgebra such that $C \neq kG(C)$ and C^* is pointed. Since U is pointed C is pointed as well. Since $C \neq kG(C)$ it follows from Taft-Wilson Theorem [TW] that C contains a non-trivial skew primitive element P . Without loss of generality we may assume that $P \in P_{a^i, 1}(U)$ for some $0 \leq i \leq N - 1$. By assumption 2, $i = \nu$. Thus $\Delta(P) = P \otimes a^\nu + 1 \otimes P$ and $k(a^\nu) \subset C$. Since P must be a linear combination of x, y and

$a^\nu - 1$ we conclude, using Lemma 2.8, that there exists an integer $1 \leq s \leq N$ such that $s|N$, $(N/s)|\nu$ and C is one of the following distinct Hopf subalgebras of U :

$$k < a^{N/s}, x, y >, \quad k < a^{N/s}, x > \quad \text{or} \quad k < a^{N/s}, y > .$$

If in addition $\omega_1^{N/s} = \omega_2^{N/s}$ then C may also equal

$$k < a^{N/s}, \alpha x + \beta y >$$

where α, β are fixed elements in k^* . Since B and H generate U it follows by the above that the mutually exclusive possibilities for B and H are the following:

1. $B = H = U$.
2. $B = k < a, x >$ and $H = k < a, y >$ (or vice versa).
3. $B = k < a, x >$ and $H = k < a, \alpha x + \beta y >$ (or vice versa) where α, β are fixed elements in k^* .
4. $B = k < a, y >$ and $H = k < a, \alpha x + \beta y >$ (or vice versa) where α, β are fixed elements in k^* .
5. $B = k < a, \alpha x + \beta y >$ and $H = k < a, \gamma x + \delta y >$ where $\alpha, \beta, \gamma, \delta$ are fixed elements in k^* , so that $\alpha\delta - \beta\gamma \neq 0$.

If 1 holds then by Proposition 3.5, $U = U_{(2\nu, \nu, \omega)}$ and we are done.

If 3-5 hold then $\omega_1 = \omega_2$. Since B^* and H^* are pointed it follows from Theorem 2.5 that for all $\alpha, \beta, \gamma, \delta \in k$ such that $\alpha\beta \neq 0$ and $\gamma\delta \neq 0$

$$k < a, \alpha x + \beta y > \cong k < a, \gamma x + \delta y > \cong H_{n_1, \omega_1, N, \nu}$$

as Hopf algebras. Therefore since $B^{*cop} \cong H$ as Hopf algebras it follows from [G2, Lemma 1.1.2] that there exists a primitive N th root of unity ω , such that $\omega^{-\nu} = \omega_1 = \omega_2 = \omega^\nu$. Therefore $N = 2\nu$, ν is odd and $xy + yx = 0$ (see the proof of Proposition 3.5). Thus $U = U_{(2\nu, \nu, \omega)}$ and we are done.

Suppose 2 holds. Since B^* and H^* are pointed it follows from Theorem 2.5 that $B = H_{n_1, \omega_1, N, \nu}$ and $H = H_{n_2, \omega_2, N, \nu}$. Thus, it follows from [G2, Lemma 1.1.2] that there exists a primitive N th root of unity ω , such that $\omega_1 = \omega^{-\nu}$ and $\omega_2 = \omega^\nu$. In particular $r_1 = r_2 = r$, so $x^r = y^r = 0$. By [R2, Theorem 2], there exists a surjection of Hopf algebras $\pi : D(B) \rightarrow U$ such that $\pi|_B = id_B$ and $\pi|_{B^{*cop}} : B^{*cop} \rightarrow H$ is an isomorphism of Hopf algebras. Let A and X be the canonical generators of B^* . Then using [G2, Lemma 1.1.2] and [R1, Theorem 4 d)] we conclude that π is given by:

$$\pi(\varepsilon \bowtie a) = a, \quad \pi(\varepsilon \bowtie x) = x, \quad \pi(A \bowtie 1) = a^m \quad \text{and} \quad \pi(X \bowtie 1) = \beta y$$

for some $\beta \in k^*$, and an integer m which satisfies $(m, N) = 1$ and $m\nu = \nu \pmod{N}$. Since the relation

$$(\varepsilon \bowtie x)(X \bowtie 1) = \omega^{\nu^2}(A^\nu \bowtie a^\nu + X \bowtie x - \varepsilon \bowtie 1)$$

holds in $D(B)$ we conclude that the relation

$$-\beta yx + \beta \omega^{-\nu^2} xy = a^{2\nu} - 1$$

holds in U . Hence after replacing $(-\beta y)$ by y we get that $U = U_{(N, \nu, \omega)}$. This completes the proof of the theorem. ■

References

- [CM] W. Chin and I. Musson, The Coradical Filtration For Quantized Enveloping Algebras, J. LMS, to appear.
- [D] V. G. Drinfel'd, Quantum groups, Proc. Int. Cong. Math. (Berkeley, 1986), 798-820, Amer. Math. Soc. Providence, RI, 1987, MR 89f:17017.
- [DT] Y. Doi and M. Takeuchi, Cleft comodule algebras for a bialgebra, Comm. Alg. **14** (1986), 801-818.
- [G1] S. Gelaki, Topics on Quasitriangular Hopf Algebras, Master's Thesis, Ben Gurion University of the Negev, Israel, 1992.
- [G2] S. Gelaki, On Pointed Ribbon Hopf Algebras, J. of Algebra **181** (1996), 760-786.
- [G3] S. Gelaki, Addendum to "On Pointed Ribbon Hopf Algebras", J. of Algebra **204** (1998), 347-352.
- [K] I. Kaplansky, Bialgebras, University of Chicago, 1975.
- [M] S. Montgomery, Hopf algebras and their actions on rings, CBMS Lecture Notes, Vol. 82, AMS, 1993.
- [R1] D. E. Radford, On Kauffman's Knot Invariants Arising from Finite-Dimensional Hopf algebras, Advances in Hopf algebras, 158, 205-266, lectures notes in pure and applied mathematics, Marcel Dekker, N. Y., 1994.
- [R2] D. E. Radford, Minimal quasitriangular Hopf algebras, J. of Algebra **157** (1993), 281-315.
- [T] E. J. Taft, The order of the antipode of a finite dimensional Hopf algebra, Proc. Nat. Acad. Science U.S.A **68** (1971), 2631-2633.
- [TW] E. J. Taft and R. L. Wilson, On antipodes in pointed Hopf algebras, J. of Algebra **29** (1974), 27-32.